

## Chapter 9

# Agroforestry: Prospects and Challenges

W. Nouman\*

### Abstract

Forest can play a vital role in improving Pakistan's economy being protective layers for watershed areas, maintaining a sustained supply of wood, wood-based and non wood forest products, conserving and improving soil fertility, erosion control, improving food, fodder and fuelwood production. As the forest area of Pakistan is very low from the standard figure i.e., only 2.5%, there is a dire need for introducing such systems and practices which can increase the forest area meeting the nation's requirements. For this, agroforestry is a better option that cannot only increase forest area but by adopting this system, marginal lands can also be brought under cultivation by incorporating salinity and drought resistant trees into land use system which can bridge the gap between demand and supply of food and fodder for livestock. This chapter describes in detail the important issues, challenges and solutions associated with agroforestry in the light of various examples, case studies and theories.

**Keywords:** Agroforestry; Multipurpose trees; Agrisilviculture; Soil improvement.

---

\*W. Nouman

Department of Forestry and Range Management, Bahauddin Zakariya University, Multan, Pakistan.  
For correspondance: wnouman@gmail.com

*Managing editors:* Iqrar Ahmad Khan and Muhammad Farooq

*Editors:* Muhammad Tahir Siddiqui and Muhammad Farrakh Nawaz  
University of Agriculture, Faisalabad, Pakistan.

## 9.1. Introduction and History of Agroforestry

Trees have an important role to play on earth by fulfilling the basic necessities of all living organisms from micro organisms to animals. The world's population is assumed to reach at 10 billion by the middle of 21<sup>st</sup> century (UN 1995) if it continues to grow at the present rate. This upcoming challenge reveals a great demand of food, fiber and shelter to cover social, medical and economical demands of human beings. Trees are also essential for the survival of about 200 million people worldwide by providing wood and income (Ansari and Iftikhar 1985).

There is no doubt that agroforestry has been practiced since millennia as an arrangement of traditional land-use systems and practices but in late 1970s, it was evolved as a modern and improved land-use system. It has been a customary practice to grow trees and crops in a combination for getting maximum benefits and intermediate income. Not only in Asia, but it has been practiced in European countries like Finland and Germany as well. Beside this, farmers of tropics and Central America have been regularly involved in growing different tropical trees on their farmlands aiming at obtaining timber benefits, shelter tree crop and windbreaks (Nair 1993). Such examples can also be found in Asia i.e., the farmers practiced agroforestry in Philippine by clearing the forest area for agricultural crop production but leaving a few selected trees to provide a partial canopy or shelter to new crop (Nair 1993). These examples indicate that trees are integral part of cropping system to support agriculture. The change of cropping pattern or the inclusion of tree species into traditional farming system started in 1806 from Burma which was under British Empire at that time. U Pan Hle, a Karen in the Tonze forests of Thararrawaddy Division in Burma, established a plantation of teak through the use of "taungya" (hill cultivation) method and presented it to Sir Dietrich Brandis (Blanford 1958). This system was further spread to other parts of Burma and later it reached South Africa in early times of 1887, India in 1890 and Bengal in 1896 (Raghavan 1960). This is important to mention that teak is not the only species that can be used in agroforestry system, other tree species of multiple benefits can also be used for this purpose.

Like the traditional land-use disciplines of agriculture and forestry, agroforestry covers biological, physical and social science disciplines. (Mercer and Miller 1998). Understandably, the biophysical sciences have dominated the first two decades of agroforestry research and development because the interest in agroforestry as a land use emerged from observations of the impacts of non sustainable farming systems on tropical soils and forests (Nair 1996). Concerns over the inadequacy of socioeconomic research in agroforestry began to grow, however, as improved agroforestry systems were transferred from research institutions to rural development projects. As mentioned above, during the 1980s agroforestry became an established focus of international rural development efforts. For example, in 1988 and 1989 ICRAF identified 166 agroforestry projects supported by developmental organizations and government agencies (Miiller and Scherr 1990), and by the early 1990s the US Agency for International Development alone supported 28 agroforestry and technological advances, agroforestry rural development efforts were frequently unsuccessful (Nair 1996).

More than a decade of discussions on how to protect the world's forests has resulted in substantial changes in the way forests are managed. Policies and programs help to promote sustainable forest management. Forest plantations comprise about 5% of the world's forests. Asia has the largest area of plantations, accounting for 62% of the world's total. China accounts for 24% of that total and India, 18% (FAO 2000). The area of forest plantations was increased by an average of 3 million hectares per year during the 1990s. Half of this increase was the result of afforestation on land previously under non-forest land use, whereas the other half resulted from conversion of natural forest. In Pakistan, at present about 90% fuel wood and 60% timber comes from farmlands. Thus, agroforestry is playing a vital role in fulfilling our wood requirements. It is estimated that 10% area of our farmlands can be easily brought under tree cover without harming agricultural crops. At present the tree cover on farmlands is only 2%. There are about 300 million trees on farmlands throughout the country with a standing volume of 70 million m<sup>3</sup> (Quraishi 1998).

The significance of wood produced on farmlands has increased sharply during the last two decades. According to FSMP (Forestry Sector Master Plan) the annual growth of forests and trees was 14.4 million m<sup>3</sup> of which 7.7 million m<sup>3</sup> (53%) was put on by the farmland trees. The farmlands of the Punjab have about 200 million trees of which 95% are in irrigated areas. These trees are mainly comprised of *Dalbergia sissoo* (42%), *Acacia modesta* (20%), *Acacia nilotica* (11%), *Melia azedarach* (7%), and Mango (6%). Ber (31%), *Acacia modesta* (20%), *Acacia nilotica* (19%) and *Dalbergia sissoo* (7%) are the predominant species in rainfed areas (Quraishi, 1998). The farmers have long recognized the value of planting trees on fields for sheltering crops, generating wood for self-consumption and commercial sale. Scattered trees have less competition with agricultural crops and they yield tangible benefits at very little cost and efforts.

Figure 9.1 manifested that agroforestry is adopted for various purposes. This figure provides a detailed overview of agriculture-forestry interface based on the socio-economic need of the community. An agroforestry system possesses three main qualities to obtain maximum benefits i.e., productivity (farm income can be increased by enhanced tree products' output, improvement in crop yield, and optimizing labour efficiency), sustainability (improvement in soil fertility, reclamation of problem soils) and adoptability (introduction of new and improved technologies in agroforestry systems).


## 9.2. Agroforestry Systems and Practices

Based upon the concept and definition of agroforestry, different agroforestry types can be observed around us. Based upon ecological, geographical and farming system combinations, different agroforestry systems can be classified. These classifications were made on the basis of findings and reports described by ICRAF and Nair (1987). These reports explain the collection and evaluation of present land-use systems. These reports not only assist in classifying the agroforestry systems but it also provided a detailed and comprehensive database to evaluate the

strengths and weaknesses of current agroforestry systems and the proposed ones. Reviewing the classification of agroforestry, a common objective can be found in all, which is the essence of agroforestry i.e., “the purposeful growing or deliberate retention of trees with crops and/or animals in interacting combinations for multiple products or benefits from the same management unit” (Nair 1993).


Generally, agroforestry systems are classified into three systems on structural basis i.e., agrisilviculture (combination of trees and crops), silvopastoral (combination of pasture, livestock and trees) and agrosilvipastoral (combination of crops, pastures, livestock and trees) (Figure 9.2). A few other classifications are also proposed based on the function, agro-ecological and environmental adoptability, and socio-economic and management level. These classification criteria are confined only to specific geographical regions. So, here we will discuss only above mentioned three agroforestry systems.

It’s a common misunderstanding that growing agriculture crops and trees on the same piece of land is considered as agroforestry. It has been wrongly interpreted somewhere in literature. For a detailed understanding about agroforestry concept, the components should be understood. In an agroforestry system, three components have prime importance i.e., trees, pastures/ livestock and agriculture crops.



**Fig 9.1** Special conditions and constraints of Agroforestry

Source: Modified from Nair (1993).



**Fig. 9.2** Classification of Agroforestry Systems

It is clear from the definition and concept of agroforestry that trees are the main components of each agroforestry system that cannot be deleted. Beside these three agroforestry systems, a few other are also in practice, for example multipurpose woodlots, apiculture (with trees), sericulture (with trees especially mulberry plants) and aquaculture/ fish farming (with trees) (Nair 1985). These systems also improve the socio-economic conditions of community livelihoods but these types do not fall in above mentioned three main agroforestry systems, hence, these are brought under *others* term. It is important to mention here that a few other sub-divisions are also in practice worldwide that include agrihorticulture (combination of agriculture crops and fruit trees) and agrisilvihorticulture (combination of agriculture crops, timber trees and fruit trees). The application of all these agroforestry systems may vary according to the main and specific objective. For example, fodder and food production, livestock production, timber production, shelterbelts, windbreaks, fuelwood production, soil rehabilitation etc.

Agroforestry practices and agroforestry systems are often used synonymously while Nair (1992) differentiate both terms as following. “An agroforestry system is a specific local example of a practice, characterized by environment, plant species and their arrangement, management, and socioeconomic functioning. An agroforestry practice denotes a distinctive arrangement of components in space and time. Although hundreds of agroforestry systems have been recorded, they all consist of about 20 distinct agroforestry practices”. In other words, in a system, more than one practice can be found simultaneously or at different times.

### **9.2.1. Trees and Arable Crops (Agrisilviculture)**

Agroforestry practices mainly consisting of trees and crops growing together are known as agrosilvicultural practices. It includes 'taungya'. In Taungya, the farmers grow their agricultural crops with young trees either on borders or in between. It's a common practice in West Africa and savannas as discussed earlier. In these areas, the farmers retain the native trees present on their farmlands and cultivate their crops amongst those trees with the addition of modern technology as hedgerow intercropping cropping or alley cropping. In such system, fast growing and nitrogen fixing tree species are preferred to retain or grow between crop lines which are periodically pruned, thinned and harvested to provide nutrient-rich mulch as fertilizer to the crop. In sloping areas, tree species are planted on contour especially to control soil erosion, improve water infiltration and reduce run-off effects. In this system, following practices have been used in various parts of the world. Some important practices in this system are elaborated here.

- Improved Fallow: In this practice, fast growing, leguminous trees are grown during a fallow period.
- Taungya: During early stages of plantation, agricultural crops are cultivated among young trees. In this practice, fast growing leguminous trees are used.
- Alley cropping: Fast growing, leguminous or fodder trees are grown between crops or as hedges.
- Multipurpose trees (MPTs) on agriculture land: MPTs Trees which can benefit in terms of timber, fuelwood, fruit and fodder are planted on bunds or terraces.
- Shelterbelts, windbreaks and live fences: tall growing trees are grown along / around agriculture land.

### **9.2.2. Trees, Livestock and Rangelands/ Pastures (Silvopastoral)**

Growing trees in combinations with pastures are termed as silvopastoral system. It includes growing and retaining forest trees in pastures to provide food and fodder to livestock and shade or shelter for both human beings and livestock. Tree browsing in silvopastoral system may occur mainly for productive or conservation purposes. For example, in United Kingdom, forest grazing in pastures is allowed periodically to conserve the habitat and retain understory ferns. Such examples have also been reported for Northern Europe, Southern Mediterranean region and Spain. Scattered trees grown or retained in dry tropical pastures contribute to feed cattle in dry seasons while in wetter tropical areas, trees may contribute in improving soil fertility by nitrogen fixation and atmospheric input.

Sometimes, agroforestry practices are encouraged without the involvement of pastures i.e., having a direct interaction between trees and livestock. In such practices, trees serve as fodder banks. Fast growing trees with better nutritional qualities are grown which can be harvested at intervals providing fodder to livestock. Moreover, trees are also used in sericulture and apiculture. These both

systems are also considered under this system but this is the opinion of a few scientists. Some common practices of this system are discussed here.

- Protein banks: Plantation of protein rich and nutritious trees on agriculture land/ pastures/ rangelands for cut-and-carry fodder production.
- Fodder Tree plantation on rangeland/ pastures: Trees are grown and retained to provide a regular supply of fodder to livestock

### 9.2.3. Trees, Crops and Livestock (Agrosilvopastoral System)

The practical approach of agrosilvopastoral system is to combine tree plantation, agriculture crop production and animal husbandry. This system is especially used to focus on conserving the resources and their efficient utilization with minimum energy losses. Here, a few main practices are mentioned which exist in this system.

- Apiculture: Trees are grown for honey production
- Sericulture: Preferably, *Morus alba* is grown for silk worm rearing.
- Multipurpose woody hedgerows: Fast growing trees with better fodder qualities are grown for livestock grazing/ browsing, mulching, fertilizer, soil conservation, etc.
- Home-gardening along livestock: Closed, multistory combination of multipurpose trees preferably fruit trees, vegetable crops around homesteads.

## 9.3. Importance of Agroforestry in Pakistan

Pakistan's economy has undergone considerable diversification over the years yet; agriculture is the most important sector of its economy. The cultivated cropped area covers 22.15 million hectares with its present contribution to GDP at 22%. Agriculture accounts for 44.8% of the total employed force of Pakistan (Government of Pakistan 2014). Most importantly 65.9% (Government of Pakistan 2014) of country's population currently living in rural areas is directly or indirectly dependent on agriculture for their livelihood. Agriculture is the largest source of country's foreign exchange by serving as base sector for its major industries viz. textile, sugar etc. Consequently, this sector has a substantial effect on the overall growth of GDP of the country. The area under cropping system in Pakistan is 22.15 million hectares (Government of Pakistan 2014) out of which 76% is cultivated area under irrigation (Government of Pakistan 2014). Being an agricultural country, mostly the inhabitants of Pakistan depend on natural resources and fulfill their requirements from agriculture, livestock, poultry, forestry and fisheries. But these resources are not too much to meet the needs of all the people, especially, those living below poverty line. It has been reported that 28.1% of rural population is living below poverty line, who are earning Rs. 878.64 per adult equivalent per month. Therefore, there is a need to justify the necessities of life by generating natural along with income resources.

Forests can also play an important role in Pakistan's economy. These are the important sources for protection of land and water resources, particularly in prolonging the lives of dams, reservoirs and the irrigation network of canals. The trees are also essential for maintaining a sustained supply of wood and wood products. Pakistan is a land of great diversity, which has yielded a variety of vegetation. According to FAO report 2000, 3.9 billion hectares or 40 million km<sup>2</sup> of the globe is covered with forests which are almost 30 percent of total land area. This corresponds to an average of 0.62 ha (6200 m<sup>2</sup>) per capita, though this is unevenly distributed. For example, 64 countries with a combined population of 2.0 billion have less than 0.1 ha of forest per capita. This estimate was based on data on forest area reported by 228 countries and territories. Among world regions, Europe (which, for the purpose of this assessment includes the Russian Federation) accounts for one-quarter of total forest area, followed by South America and then North and Central America. South America is the region with the highest percentage of forest cover (almost half of the land area) and Asia is the region with the lowest percentage of forest cover (less than 20% of land area). If a comparison is carried out among Asian countries, India stands first representing 21.6% of its area under forest and China ranked 2nd having 17.5% while Pakistan has only 3.1% or about 1,902,000 hectares of total land area while in 2005 it remained only 2.13%, which is very low as compared other Asian countries (World Bank 2011). The figure of Government of Pakistan is different from above. She reported that the area under forest cover in Pakistan is about 5%, out of which 85% is public forest, including 40 % coniferous and scrub forests on the Northern hills and mountains (Government of Pakistan 2005). The balance is made up of irrigated plantations and Riverain forests along major rivers on the Indus plains, mangrove forests on the Indus delta and trees planted on farmlands. If we consider the official figure of Pakistan, then it is also less than the world forest area, which is 30% of total land. Forest cover is divided into four percentage groups (>70%, 40-69%, 10-39%, 0-9%), thus the Pakistan lies in the last category i.e., 0-9%. Moreover, it can further be categorized into primary, protected, conserved and production forests (FAO 2000).

Trees are being cut at in an unmanaged system i.e., there is no balance between harvesting of trees and regeneration or afforestation activities. Such brutal practice is fashioned due to increasing demands for fuel to meet domestic and industrial needs. Unbalanced harvesting and afforestation activities hamper forest decline rate which is consequently leading towards environmental and land degradation, air pollution, and biodiversity loss. For this, efforts should be undertaken to multipurpose tree plants new and improved varieties of trees and plants that have a good growth rate. It has been reported that Pakistan lost about 214,000 ha forest @ 2.1% deforestation rate only between 2000-2005 which correspond to 43,000 ha/year loss. Pakistan's total deforestation rate from 1990-2000 was 41,000 ha or 1.8% per year. Hence this report shows that the deforestation rate in Pakistan is increasing day by day. This is very alarming figure of a great concern (FAO 2000).

Pakistan is forest deficient country with only 0.03 ha as compared to world average of 1 ha per capita (Government of Pakistan 1992), is facing timber and fire wood shortage of about 29 million cubic meters (Government of Pakistan 2005) and our

forests produce only 32% of fuel wood supply to meet national energy requirements (Government of Pakistan 1992). In a study, it has also been estimated that this shortage of 29 million cubic meters will become 52.6 million cubic meters up to 2018 thus increasing future needs by 23 million cubic meters (Wani 2003). This forest area per capita is declining due to growing population rate at 1.90% annually. According to safe estimates, the annual import bill for pulp, paper and paper products runs into RS 8 billion which is expected to increase many times due to increase in population. The area under public forest cannot be further expanded to keep pace with population growth rate and increasing demands for forest products. The only available option is to increase wood production on private or farmlands so as to meet pulp and paper demand locally to reduce import bill and save foreign exchange. It is estimated that state forests contribute only 14% of timber and 10% of fuel wood whereas 46% of timber and 90% of fuel wood requirements are being met from farmlands (Government of Pakistan 2014). To counter the deficit issues in timber and fuelwood supply, agroforestry is an important practice. It will be discussed in next sections of this chapter that how agroforestry can be helpful in soil conservation, improving soil fertility, erosion control, for improving food, fodder and fuelwood production of the country. Below are some facts demonstrating the role of agroforestry in developed and developing countries.

- Intercropping of Paulownia elongate with cereals is practiced on 3 million hectares in China (Sen 1991).
- In Henan province of China, agroforestry was introduced in late 1970s. 30 years after initiation of agroforestry practices, two-thirds of the 46,000 ha of farmland were intercropped with Paulowina elongate (Wu and Zhu 1997).
- In Tabora District of Tanzania, about 1,000 farmers intercropped *Acacia crassicarpa* with tobacco and maize (Ramadhani et al. 2002).
- In Uttar Pradesh, India, 30,000 farmers cultivate *Populus deltoids* on their farms. The average farm size under populous deltoids plantation is 1.3 ha. These plantations support their national match industry and minimize the import cost. (Jain and Singh 2000; Scherr 2004).
- According to McAdam et al. (1999) ash trees were intercropped with ryegrass pastures at 40 years rotation in United Kingdom. No reduction in ryegrass yield was recorded for first 10 years of rotation.
- In the Canada, more than 43,000 km of windbreaks has been planted since 1937 which protect 700,000 ha of Canadian prairies.
- In 1987, approximately 858,000 windbreaks were planted in the United States, especially in the north central and Great Plain areas, covering 281,000 km which protected 546,000 ha (Williams et al. 1997).
- An increase of 8, 12, 23 and 25% in yield of spring wheat, maize, winter wheat and barley, respectively was recorded by Kort (1988).

## 9.4. Agroforestry on Marginal Lands

Poor physical and chemical quality of soils severely affects the crop growth and productivity (Flowers 2004). A vast area of the world is wasteland due to excessive salts, waterlogging, acidity, alkalinity, or even the presence of sand or gravel in soil (Zhu et al. 1997). Agroforestry practices can be implemented to bring such wasteland under cultivation by imparting different techniques (Munns 2002; Munns and Tester 2008). In general, selection of suitable tree species especially multipurpose trees is the key factor in reclaiming such kind of areas (Cavalcante and Perez 1995). These trees not only provide timber and fuelwood to the farmers but rehabilitate the soil conditions improving its fertility. Detailed discussion can be seen in Chapter No. 14.

## 9.5. Agroforestry for Soil Conservation

Agroforestry is a unique, ecologically balanced and sustainable ecosystem. This is greatly to the fact that trees being a woody component offer the most: food, fuel wood, fodder, shade etc. They also minimize nutrient losses as a result of leaching and erosion. The large tap root with wide spreading lateral roots not just provide anchorage but also play an important role in nutrient uptake, absorbing nutrients from the lower strata of the soil. Nutrient cycling is another major aspect whereby the nutrients absorbed by the tree are returned back in form of litter fall, stem flow and decomposition. This again provides nutrients for fresh use to the same trees. Soil microbial associations such as rhizobia with the tree roots improve nutrient uptake especially in the case of nitrogen. Such fast growing species combined with herbaceous crops enhance the soil fertility status and improve the microclimate, improving survival conditions for both plants and animals. The roots of the trees are long to give the tree anchorage and wide to avoid soil erosion. Some trees fix nitrogen and produce nitrates which are essential for plants. The mycorrhizal roots play a vital role in the poor and the dehydrated soil. Productivity and protection of system and soil conservation are certainly most important functions of trees. Though originally equated with control of erosion, soil conservation now associates with maintenance of soil fertility (Young 1987). Moreover, the plantation of trees improves physical and chemical characteristics of soil by adding soil nutrients through leaf litter decomposition. Tree roots uptake soil nutrients below crops' roots level and make them available by fixation (Hartemink et al. 1996; Allen et al. 2004). In this way, trees can maintain soil organic matter and nutrients but it depends on tree species, planting space, tree age and agroforestry system (Szott et al. 1991; Mohsin et al. 1996; Young 1997). For example, *Populus deltoides* is considered as a potential multipurpose tree or agroforestry system as it has single and tall stem, fast growth, deep roots and narrow crown which enable less competition with associated crops (Jain and Singh 2000).

The researchers have mentioned certain qualities which are necessary for trees to maintain soil status like a high yield, better aboveground biomass production, more nitrogen fixation, dense adventitious roots capable of mycorrhizal association, existence of deep roots, less toxic substances and adequate amount of nutrients.

There are many mechanisms involved in soil fertility improvement such as compatibility of trees and their suitability to the soil. A leguminous tree fixes nitrogen and improves nutrient status. Leaf litter increases organic content. The organic matter is greatly influenced by the microbial activity in the rhizosphere, which are beneficial for the companion crops. Physical properties such as water holding capacity (WHC), permeability, aggregate stability of soil and soil temperature are some of the more essential aspects. Soil conservation and erosion control through biological measures lend stability to the system and decrease siltation rate in downstream reservoirs.

Previous research reports, observations and farmers' experiences support the hypothesis that the plants improve the soil quality, texture, organic matter and nutrients accumulation. Following points strongly support this concept:

- From ancient times, the farmers have been interested in cultivating their agricultural crops on cleared forest lands.
- Forest soils have been found with better physical and chemical properties pertaining ample amount of organic matter and essential nutrients in comparison with fallow lands.
- Being a closed ecosystem, woodlands are preferred for nutrient transfer, storage, and cycling.
- The best way to reclaim a marginal land or improve the soil fertility on sustainable basis is to incorporate tree cultivation either as compact plantation, linear or alley cropping. It will gradually improve soil organic matter, nutrients accumulation and water holding capacity.

The summarized beneficial effects of agroforestry on soil fertility and productivity are given in Table 9.1.

Nutrient cycling includes the conservation of sources and their partitioning between soil and plant within a system. Agroforestry systems have been cited as potential land-use systems for their efficient nutrient cycling which in turn improve soil fertility. As discussed earlier, trees possess deeper root system than other herbaceous plants having the potential to retrieve nutrients from deep soil horizons. Moreover, forest, an undisturbed ecosystem, takes nutrients from the atmosphere, or by nitrogen fixation etc enabling the atmospheric input to soil. Such input of biologically important elements is caused through precipitation. Nitrogen is essential element as first floor contains 2225 kg/ha which is mostly in humus complexes (Young 1987). The amount of nitrogen and potassium is higher in the leaves and fruit of plant than in branches, poles and litter. Leaf fall is an important process as the litter continuously provides the nutrients back to the soil and they keep the cycle in progress. With respect to the nutrient intake, the major part of nutrients is in the leaves of the nutrients. The mechanisms include uptake from lower soil horizon, reduction of leaching to lower horizons, balanced nutrient supply and improvement ratio between available and fixed nutrients for a tree leave biomass production of 4000kg dry matter per ha year. The potential nutrients returns in form of nutrients are of the order of 80-120 kg N, 8-12 kg phosphorus, 40-120 kg potassium, and 20-60 kg calcium. The nutrient uptake in trees is currently

explained by a hypothesis that trees are most efficient agents than the herbaceous plants in intake of nutrients. Potassium, phosphorus, bases and macronutrients are released by rock weathering particularly in the B/C and C-horizon on to which tree roots often penetrate. The strong gradient in nutrient content, control between top soils and sub soils indicate recycling through litter whereas other processes are also involved. It is important to note here that almost 20-30% of tree total living biomass is present in soil as roots which are constantly adding organic matter to soil through its dead and decaying parts.

**Table 9.1** Beneficial effects of trees on soil fertility

Addition to soil	Reduction of losses from soil	Effect on physical properties of soil	Effect on chemical properties of the soil
Sustainable supply of organic matter to soil	Protection from soil erosion: proper planning and management of trees on wind/ water eroded soil can minimize the effects of soil erosion in terms of loss in organic matter	Improvement of soil physical properties i.e., soil structure, texture, porosity, water holding capacity	Decrease in soil acidity
Nitrogen fixation through leguminous tree roots	Nutrient reclamation: soil nutrients which are deeply deposited in soil horizons are absorbed and uptaken by tree roots	Amelioration of soil temperature	Decrease in salinity and sodicity
Nutrient uptake as trees are more active in uptaking soil nutrients from deep root zones especially in B/C and C horizons Atmospheric input: nutrients dissolved in rainfall and then deposited in soil			Effect of shading: by lowering down soil temperature, loss of organic matter through oxidation can be minimized

The plantation of trees in the agroforestry systems increases soil fertility by improving the physical properties of the soil. Sreemannarayana et al. (1994) studied influence of *Albizialebbeck* on soil properties with cowpea as intercrop in different seasons and with pruning treatments. The soil organic matter improved over the recorded value. Great improvement in soil fertility was shown by *Albizia lebbeck*. Karite (*Vetellaria paradoxa*) and Ner (*Parkia bilobosa*) were observed in this regard. They showed that the organic matter was rich in the soil under the trees and soil fertility improvement was least in the fallow land and was highest in *Albezia lebbeck* based system. Induced crop yield are from 0-60%. The study in Rampur (Utter perdes, India) shows benefit of poplar (*Populus deltoides*) plantation. The

litter produced had higher content of N P K. Alley cropping has also proved much beneficial. Soil carbon was found to be 8.7 % higher than sole crop plant. However, C/N ratio decreased.

When woody perennials and agricultural crops are together cultivated they improve soil conditions and fertility greatly. The productivity, fertility, and erosion under cassava based agroforestry systems were studied by Gosh et al. (1989). There was increase in the phosphorus and potassium status. Alley cropping also has a significant effect on it. It showed that large biomass could be obtained from this system. In most cases 30% of nitrogen is available; the rest is lost by leaching and gaseous losses Gosh et al. (1989). In leucaena, 50% of N is released in first 25 days.

The trees may have negative effects on the crops and soil fertility: if the tree plantation is poorly managed it leads to soil erosion. Similarly shading and spectral quality of light in close proximity may have serious effects on agroforestry and adverse chemical and biological effects may result from acidification, allelopathy, etc. According to a phenomenon, the tree roots and leaves have some substances that hinder crop production regardless of their benefits like benzoic and *p*-coumaric acids can create a problem if they are present in large quantity in the trees.

Application of fertilizers to forest trees in agroforestry is generally a costly process but it suits to the trees of economic values as seen in horticulture based trees. An ideal fertilizer must be cheaper to get a unit of plant food, give high growth response, maintain fertility and physical and chemical properties of soil (Maslekar 1990). Generally, hardwoods require more nutrients than conifers and fast growing trees.

Many advanced countries except India have accepted the benefits of fertilizers and it is proved to be better for getting large biomass in short span of time. Significant growth advancements due to the putting of 28 kg N and 15 kg P<sub>2</sub>O<sub>5</sub> per ha were observed in pines (Singh et al. 1981). Krohn (1981) reported that fertilization of eucalyptus seedlings resulted in better tree growth and survival. The effect of phosphoric fertilizers on growth of teak plantation was also beneficial. Ammonium phosphate also proved beneficial for growth and height of teak plants in initial 2 years. In case of SS, a dose of 120 g per plant showed significant effect on height (Singh et al. 1981)

Generally, soils brought under agroforestry systems are in poor condition with low nutrient availability, and mismanagement may lead to mortality of trees soon after planting. Supplementary nutrients should be ensured at the critical stage of planting. Fast growing leguminous species ensure increased fertility of soil along with herbaceous cover as this constantly contributes in biomass and organic matter in form of leaf litter. This increases nutritive status and prevents erosion. Any biomass that accumulates is decomposed and liberates nutrients for reuse. Further techniques can be introduced for efficient recycling, soil aspects should be assessed in detail. Any negative effects should be remedied in accordance with expert advice and dealt with immediately.

## 9.6. Interventions for Soil and Water Conservation Through Agroforestry System

Soil and water are the most important resources for life. However, ignorance of soil and water conservation measures, improper use of fertilizers, mismanaged irrigation and water practices, mining, and effluents discharge etc. are responsible for land degradation at a massive scale. This leads to the paradox of getting more production from the increasingly scarce resources to meet the requirements of mounting human population. Demographic pressure is the primary cause of land degradation, along with loss in vegetative cover. Shifting cultivated areas may result in further losses of 18 million tonnes of soil. Values for tributary river sediments in the region range from 6.0 to 98.4 cum ha<sup>-1</sup> per year. Hence agroforestry techniques become crucial for the conservation of soil and water.

Water erosion can cause losses of top soil in 130.5 million ha as well as degradation of terrain in 16.4 million ha. Of the total eroded soil, 29% is lost to sea, 61% relocated or transferred to different places, and 10% is sedimented in reservoirs, reducing their capacity by 1-2 % at an annual basis. The erosion rates have reduced the life of reservoirs increasing rates of sedimentation, especially in the catchment areas.

If we talk about creating barriers for controlling run-off and soil loss on slopes, the most effective and feasible are the vegetative barriers. *Panicum maxicum* used as a barrier in maize succeeded in decreasing run off by 28% and erosion by 48% at Dehradun (Bhardwaj and Kholia 1999). The barriers also help in redeposition of soil on contours forming bunds or terraces naturally. However, the effectiveness of the barrier also varies according to situation and conditions. For example, in Doon valley run-off was reduced by 18% and soil loss controlled upto 78% on a 4% slope using barriers of native grasses such as *Vetiveria zizanioides* (Vetiver), *Eulaliopsis binata* (Bhabar) and *Panicum maxicum* (Guinea). But the same grasses cannot be universally accepted to achieve the same result or control erosion and run-off in different countries or zones.

Apart from grasses, tree species too prove to be an effective control measure. Paired contours and hedgerows consisting of *Leucaena leucocaphala* (ipil ipil) and Eucalyptus hybrid (Sufeda) can reduce run-off by 40-48% as well as 12.5t ha<sup>-1</sup> soil loss in maize on 4% slope. Other suitable grass species include *Panicum maxicum*, *Chrysopogan fulvus*, *Setaria sphacelata*, *Saccharum munja*, *Saccharum spontaneum* and *Arundo donax*. *Panicum maxicum* reduced soil losses from 45t ha<sup>-1</sup> to 6.12t ha<sup>-1</sup> in Dehradun (Bhardwaj 1990).

Hedgerows are of multipurpose concept. They don't just reduce soil loss and run-off, but provide fodder too. They also trap soil particles helping to prevent loss of nutrients and finer soil particles. Even small farms can easily raise hedgerows. Grewal (1993) compared agroforestry models for eroded lands stated soil loss to be most negligible when under a Eucalyptus hybrid and *Eulaliopsis binata* system, soil loss being only 0.07t per ha. *Acacia catechu-Pennisetum purpureum* was next with loss of 0.24t per ha, and *Tectona grandis-Leucaena leucocephala-Eulaliopsis*

*binata* with loss of 0.43t per ha. Run-off and nutrient losses were reduced too. Eucalyptus trees managed as a sole tree are more effective. Grass cover decreases run off by 73 percent and soil loss by 94 percent. These were further reduced by eucalyptus tree rows. The trees affect the crop yield but this is compensated by tree products (Narain et al. 1998). Local flash floods can be controlled by establishment of vegetative spurs, brushwood check dams, vegetative filters and stream bank lining of native species. This will save a greater portion of land from degradation and erosion (Samra 1994). Leguminous shade trees are of immense importance to tea plants as they provide shade and nitrogen (Tejwani 1994). Shade trees provide 2500-5000 kg leaf litter, 63-126kg N, 18-36kg P<sub>2</sub>O<sub>5</sub>, 22-44kg K<sub>2</sub>O, 32-64kg CaO and 16-32kg MgO ha<sup>-1</sup> per year to soil (Gogoi 1976). These trees play a vital role in improving fertility and enhancing production by improving soil nutrient content.

Mine spoils, or mining, results in degradation of land by shifting of soil, denudation, reduction in water holding capacity of soils etc. The disturbed land can be reclaimed by establishing forest cover with trees as well as grasses and shrubs. Agroforestry is the ultimate option for such a solution. Species that are indigenous, hardy, with coverage and nitrogen fixation with good economic and social values are preferred. From the western states of Jammu and Kashmir to Himachal Pradesh, about 10 million ha of land are covered by the cold desert as it lies in the Himalayas and Tibetan plateau. (Rai 1996) lying in the rain shadow of mountains and facing severe extremes of cold and dryness, this is a fragile and unstable area. The constant biotic pressure in terms of grazing, fodder and fuel has led to degradation. Faulty land practices and exploitation of tree cover are other reasons. Eucalyptus and poplar clones have been recently introduced in the Kashmir Valley. Other agroforestry species suited to the region are *Quercus* spp., *Bauhinia variegata*, *Morus cerata*, *Ulmus wallichiana*, *Celtis australis* etc. A native shrub, *Hippophae rhamnoides*, is used for both fodder and fuel with promising potential. One ha plantation of the shrub is sufficient to fulfill needs of 20 households. (ICFRE 1993)

Integrated watershed management is an essential to reduce run-off, flooding in down-stream area and improved in-situ moisture conservation, hence improving biomass production (Samra 1994). Vegetation in the catchment area helps ground water recharge and rise in water table. Both run-off and soil loss are decreased when watershed areas are incorporated with trees and grasses, or mechanical measures are taken. (Singh et al. 1990)

To conclude, agroforestry has the distinctive feature that the land use systems have the potential for both production and conservation. (Young 1986) These include products and services such as conservation of resources, prevention of erosion, increased soil fertility, shade, moisture conservation, recreation, biodiversity, etc. The agroforestry interventions are aimed to increase biomass production, soil amelioration and stabilization of degraded lands. Not only are the needs of fodder, food, fuel wood, fiber satisfied but economic and environmental security is ensured too, leading to sustainable development with steady income sources.

## 9.7. Agroforestry for Food Production

Development and well-being of humanity depends on the degree of sustainability of human activities. A sustainable community must be able to balance the socio-economic development with the production of food and the environment protection. The speedily inflating human population is pushing for greater food requirements, which is in turn, threatening the environment conservation and enlarging the gap between the supply of resources and the demands of basic necessities of life. Since ancient times, people are fulfilling their requirements for food and shelter by depleting natural resources. The scientists and policy makers are encouraging communities to adopt sustainable ways of harvesting benefits from natural resources without causing any serious environmental concern. Fischer et al. (2009) reported that 12% of total world population (800 million) is undernourished and the policy makers are devising protocols to minimize this figure to its half by 2015. The food security issues can be resolved by imparting various agricultural approaches and technological innovations in current farming system. Currently, a wide array of fruit trees, crops, vegetables, etc. are being practised in agricultural system. However, food shortage threats can be overcome by planting under-utilized forest trees rich in minerals, protein, fiber, etc. (Leakey 1999; FAO 2010; Malézieux 2013). The scientists are working on the domestication of forest trees to provide year around income, food for local communities and feed for livestock. For this, fruit trees are being utilized as primary option. This said, the significance of wood-based and non-wood based forest products cannot be declined either. A case study in Africa revealed that 90% households cultivate fruit trees. Of these, one fourth of the respondents grew banana, avocado and mango while two third were growing some other fruit species. Likewise, the people of Malawi consume papaya and oranges which are mostly harvested on their own farmlands (Jamnadas et al. 2011).

In addition to the provision of food, feed, fruits and nutrients, forest trees offer meaningful aid in sheltering, nourishing and improving agricultural crop systems through means like providing shade for crops that are sciophytic in nature (grows well under shade conditions); shelter and fodder for livestock production and improving soil fertility through litter-fall and extensive adventitious and tap root system (*how tree root system benefits soil fertility and crop productivity is discussed in "Role of Agroforestry in Soil Fertility"*). Nitrogen fixing trees improve crop yield by improving soil fertility, rain use efficiency of crop plants and better performance under drought conditions (Sileshi et al. 2011). Moreover, these tree species can regulate the climatic factors especially micro ones which can assist agricultural crops to mitigate harsh climatic effects resulting in better growth behaviour and higher yields (Sileshi et al. 2012). In addition to supporting agricultural crops, trees also play their roles in dairy farming by providing, shade, shelter and fodder and supplementary feed.

Home gardening has been considered as a viable option to support food requirements of households. Home gardens are preferably a combination of plants, shade trees, fruit bearing trees, shrubs, vines and medicinal plants. In home gardens, tuber crops, medicinal plants and fruit trees are preferred crops since these

can fulfil the basic needs however, a conspicuous character of a home garden is the dominance of fruit trees like mango, guava, date palm etc. and food providing trees like *Moringa oleifera* and *Sesbania grandiflora* which dominate in Asian home gardens. These trees make up the deficiency of local diet providing better nutritious supplements. It has been reported in literature that home gardens support the energy and food requirements of local farming communities substituting vitamins and nutrients' deficiency (Soemarwoto and Conway 1991).

In order to increase the function of agroforestry in food and nutritional security, following recommendations may be considered:

- The present role of agroforestry tree products and services in sustaining food and nutritional security of inhabitants of rural area in different farming systems should be better quantified.
- The expansion of agroforestry policy should not be restricted to the agricultural or forest sectors only. It should be treated as an individual farming system. Required improvements include selecting suitable tree species and land occupancy, in-depth research on practices and behavior of farmers to grow these trees and the credit of agroforestry as an investment option for food production.
- Research studies should be focused on investigating food tree domestication options appropriate for meeting smallholders' needs, and assess complementarity and resilience in agroforestry systems under climate change in the context of other global challenges to food and nutritional security.

## 9.8. Agroforestry for Fuelwood Production

Traditional energy sources could not have much consideration in current energy debates, but firewood and charcoal from trees are vital for the endurance and well-being of approximately two billion people (FAO 2008). In sub-Saharan Africa, the use of charcoal is increasing speedily, as the value of charcoal industry is supposed to reach about US\$ 8 billion in 2007 (World Bank 2011). The charcoal industry is an essential element for food and nutritional safety because it provides both energy and income. As the prices of modern energy sources are increasing, this situation is improbable to change for some time.

In developing communities/ in below-the-poverty-line/In communities with inadequate economic means, firewood and charcoal are frequently burnt in open fires or poorly-functioning stoves with considerable release of pollutants (especially from firewood) that may be fatal to human health. Such a condition has already lead to the deaths of more than one million people in a year worldwide. Of these, women stand most victimized since they are in direct contact with cooking activities (Jamnadas et al. 2011). Different tree species exhibit varying fuel quality as mentioned in "Multipurpose Trees MPTs".

Reduced access and increased prices of wood-based biomass have led to initiatives to promote agroforestry cultivation. The farmers who are involved in agroforestry practices growing their own fuelwood trees need less fuelwood to purchase from market as most of their needs can be fulfilled from their own sources. Moreover, there is less reliance on natural forest stands and requires lesser time in collection. This leaves more time for income-generating activities, especially for women who are usually the major fuelwood collectors (Thorlakson and Neufeldt 2012). Access to cooking fuel provides people with more flexibility in what they can eat, including foods with better nutritional profiles that require more energy to cook. The cultivation of woodlots allows the production of wood that is less harmful when burnt and has higher energy content. The use of better stoves – with greater efficiency – reduces greenhouse gas emissions relative to the energy generated for cooking purposes.

Pakistan is an energy deficient country. It has been reported that approximately 54% of national energy demand are covered by electricity conventional energy sources while the utilization of fuelwood, crop residues and dung fulfill the remaining requirements. It is important to mention here that fuel wood trees fulfill 26% of total energy requirements. By now, 95% energy requirements in rural areas are fulfilled by trees while in urban areas, this figure goes to 56% (Zaigham 2004). According to Zaman and Ahmad (2012), total wood consumption of Pakistan would be 59.44 million m<sup>3</sup> including timber consumption (16.62 million m<sup>3</sup>) and fuelwood consumption (42.81 million m<sup>3</sup>) by 2025 if population is 208.84 million. Currently, per capita fuelwood consumption is estimated at 0.205 m<sup>3</sup> while total fuelwood consumption is 34.95 million m<sup>3</sup> in Pakistan. Three main sectors are fuelwood consumers i.e., household, commercial (restaurants, hotels, tea shops, bakeris, milk shops etc. and industrial sector (mainly brick and tobacco curing industries). Zaman and Ahmad (2012) reported household sector as the largest fuelwood consumer (81.8%) followed by industrial fuelwood entrepreneurs (14.9%) and commercial sector (3.3%). Out of household sector, 75% households consume fuelwood for cooking purposes, 14% for water heating while the rest (11%) for room heating in winter. Moreover, rural community consume a major part of fuelwood for cooking purpose (90%) while in urban areas, only 10% inhabitants depend on fuelwood (Government of Pakistan 2011). In case of fuelwood consumption by industrial sector, major proportion goes to social ceremonies (27%), khoya production (24%), brick making (20%), charcoal making (8%), tobacco curing (3%) and other industries (18%).

Mainly *Acacia nilotica*(kikar), *Acacia modesta* (phulai), *Dalbergia sisso* (shisham), *Ziziphus jayuba* (ber), *Morus alba* (mulberry) are used for fuelwood consumption depending upon their calorific values, physical and chemical characteristics. Tree species with minimum moisture percentage are preferred that produce more heat as compared to other species with higher moisture content.

The selection of suitable tree species plays a key role in the production and exploitation of fuelwood in an agroforestry system. The selection mainly depends on community preferences, climatic factors, edaphic factors, landholding size, interaction with existing agricultural crops, and market demands. So, it is rather

difficult to propose a specific fuelwood tree which can address global requirement but a few criteria can be made to select suitable species.

- Adaptation ability to local environmental conditions. In this case, indigenous species are preferred which are of community interest.
- Continuous pruning can supply unremitting fuelwood but for this, highly branched tree species should be preferred which can endure pollarding and pruning shocks.
- Multipurpose trees provide a number of useful benefits like branches can be used as fuelwood and leaves can be used as livestock fodder etc.
- Spineless wood of small diameter is easy to harvest and transport because that can be easily utilized for cooking and heating purposes.
- TCI (tree-crop interaction) is an important factor while selecting a suitable tree. The species which are compatible with agricultural crops should be preferred.
- Species with low moisture contents, higher density, and more ash and sulphur content possess good combustion quality.
- The consumers' demand is highly influential in selection.

A large number of tree species have been identified as fuelwood crops. Fuelwood markets are dominated by such tree species as *Acacia nilotica*, *Tamarindus indica*, *Prosopis cineria*, *Salvadora oliodes* and other local species, inspite of the large scale tree planting efforts for fuelwood production by state agencies using exotics such as *Leucaena*, *Casuarina* and *Eucalyptus*.

Increasing efforts are being made all over the world to plant new species and improved varieties of trees that have good growth rates as well as have multiple uses. In Pakistan, many efforts are being made for introducing and cultivating fast growing and short rotation trees. The species most favored by farmers are kikar and shisham. They use the wood of these tree species for fuel, shelter, shade and fodder.

## 9.9. Multipurpose Trees (MPTs)

Multipurpose trees are defined for their multiple benefits which can be driven from their different parts like food, fruits, nuts, fodder, timber medicines etc. This term is commonly used in agroforestry especially when the farmers are to decide about selection of suitable tree species. No doubt, all trees provide several benefits like soil improvement, shade, shelter etc. but MPTs have greater impact on farming culture because trees are planted not only for one purpose. The farmers are willing to get fodder, timber and fuelwood while planting trees as living fences, shelterbelts or alley cropping. In this part, scientific and local names, family and sub-family, origin, distribution, habitat and ecology of potential trees which can be used in agroforestry systems are given in this table with their uses, specific gravity, calorific values and productivity (Table 9.2).

**Table 9.2.** Multipurpose Trees for Agroforestry

No.	Scientific name	Local name	Family-Subfamily	Origin	Distribution	Habitat & Ecology	Uses	Specific Gravity	Calorific Value	Productivity
1	Acacia catechu	Katha	Leguminosae-Mimosoideae	Subcontinent	Malakand, Hazara, Rawalpindi, and plain areas of Punjab	Drought tolerant, grows best on rocky, stony, sandy and loamy soils, required precipitation is 500-2700 mm/year, upto 1200 m elevation, -5 to 40 °C	Fodder, food, agriculture implements, handy tools, timber	1	5200 kcal/kg	4-7 m <sup>3</sup> /ha/yr
2	Acacia nilotica	Kilker	Leguminosae-Mimosoideae	Pakistan	Sindh, Punjab, Balochistan, KPK	Drought and salinity tolerant, requires precipitation 125-1300 mm/year, semi arid, subtropical/tropical climate, 1-45 °C	Fodder, fuel, charcoal, agricultural implements, land stabilization, nitrogen fixing	0.75	4900 kcal/kg	4-15 m <sup>3</sup> /ha/yr
3	Acacia senegal	Gum Arabic	Leguminosae-Mimosoideae	Pakistan	Sindh and Balochistan	Drought resistant, usually below 1700 m elevation, intolerant to waterlogging, requires precipitation between 200-800 mm/year, arid to semi-arid, hot subtropical climate, -4-48 °C	Fodder, fuel, charcoal, agricultural implements, land stabilization, nitrogen fixing	3200	kcal/kg	1-4 m <sup>3</sup> /ha/yr
4	Albizia lebbek	Kala Sirin, Black Sirin	Leguminosae-Mimosoideae	Sub-Himalayan tract	Sialkot to Hazara, Bajor, Buner, Malakand and plains of sindh & Punjab.	Well drained and loamy soils, pH 8.7 to 9.4, Summer precipitation of 400-1000mm/yr. sub-humid/sub-tropical/tropical climate, 4-40 °C, elevation 0-1600 m.	Fodder, fuel, agriculture, agricultural implements, land stabilization	0.55-0.64	5100 kcal/kg	5 m <sup>3</sup> /ha/yr

No. Scientific name	Local name	Family-Subfamily	Origin	Distribution	Habitat & Ecology	Uses	Specific Gravity	Calorific Value	Productivity
5 <i>Albizia procera</i>	Sufed Sirin, White Siris	Leguminosae- Mimosoideae	Central/South India, Bangladesh, Burma	Punjab, KPK	Prefers moist sites, tolerant to saline/sodic conditions, summer precipitation of 500-1000mm/yr, subhumid/warm/subtropical climate, temperature 1-45 °C, elevation 0-1200 m.	Fodder, fuel, apiculture, agricultural implements, tannin, furniture, poles & construction.	0.69	4800 kcal/kg	10 m <sup>3</sup> /ha/yr
6 <i>Azadirachta indica</i>	Neem, margosa tree	Meliaceae	India, Pakistan, Nepal, burma, Afghanistan, China & Sri Lanka	Sindh, Southern Punjab, Lower Balochistan	Rich loams to nutrient deficient soils that are not saline or waterlogged (water table > 18 m), precipitation of 300-1150 mm/yr, arid/hot tropical/subtropical climate, 1-45 °C, prone to frost damage.	Furniture, fodder, wood carving, medicinal, oil, tannin, timber and agricultural implements.	0.68	4990 kcal/kg	5-18 m <sup>3</sup> /ha/yr
7 <i>Bombax cibeia</i>	Simal, Silk Cotton Tree	Bombaceae	Pakistan, India, Nepal	Sub- Himalayan tracts and eastwards from Hazara.	Best in well-drained deep alluvial soils, precipitation 750-1700 mm/yr, humid warm subtropical/tropical monsoon climate, -5 to 40 °C, elevation of 1000 m, drought hardy but not frost tolerant..	Fuel, ornamental, medicinal, canoes, furniture, carvings, seed cotton for pillows.	4900 kcal/kg	Diameter growth: 35 cm/yr	

No. Scientific name	Local name	Family-Subfamily	Origin	Distribution	Habitat & Ecology	Uses	Specific Gravity	Calorific Value	Productivity
8	<i>Cassia fistula</i> Amaltas, Indian Laburnum	Leguminosae- Caesalpinoideae	Pakistan	East of Indus, upwards to Himalayas, and throughout plains.	Moderately shade tolerant, grows on variety of soils, sub-humid cool to warm/tropical humid precipitation of 500-3000 mm/yr, -5 to 45 °C, prone to frost damage.	Fuel, ornamental, agricultural implements, fine furniture, tool handles, support posts, cart wheels, tannin, medicines	5164	5164 kcal/kg	10-12 m <sup>3</sup> /ha/yr
9	<i>Dalbergis sissoo</i> Shisham, Tahli, Rose wood	Leguminosae- Papilionoideae	Sub-Himalayan tract of the subcontinent	Mostly along river banks and streams, in plains and foothills.	Frost hardy, dry subtropical/dry temperate climate, well drained sandy/sandy loam soils, elevation 900-1500 m, precipitation of 300-2000 mm/yr, temperature range of 0 to 50 °C	Fodder, furniture, fuel, charcoal, railway carriages, sports goods, farm implements and medicinal.	0.85	5000 kcal/kg	7.7 m <sup>3</sup> /ha/yr
10	<i>Eucalyptus camaldulensis</i> Sufeda, Lachi, Red River Gum	Myrtaceae	Australia	Plains and hills.	Intolerant, frost hardy and may even tolerate drought, variety of soils even saline/sodic/waterlogged, semi-arid/warm hot/subtropical winter/monsoon climate, temperature -5 to 40 °C, elevation of upto 1400m, precipitation of 200-1250 mm/yr.	Carriages, fuel, furniture, oil, charcoal, apiculture, shelterbelt, pulp & fiber board	0.71	4900 kcal/kg	10-25 m <sup>3</sup> /ha/yr

No. Scientific name	Local name	Family-Subfamily	Origin	Distribution	Habitat & Ecology	Uses	Specific Gravity	Calorific Value	Productivity
11 <i>Eucalyptus citriodora</i>	Sufeda, Lemon Scented Gum	Myrtaceae	Australia	Plains and hills.	Well drained soils, tolerates dry periods for 5-6months, semi-arid/warm hot/sub-tropical winter/monsoon, precipitation of 600-900 mm/yr, 5-40 °C, elevation upto 2000 m, tolerates light frost, easily coppiced	Fuel, charcoal, perfume, furniture, shelterbelt, apiculture, fiber board, pulp and tool handles.	0.78	4800 kcal/kg	10-15 m <sup>3</sup> /ha/yr
12 <i>Eucalyptus microtheca</i>	Sufeda, Flooded Box	Myrtaceae	Australia	Plains and hills.	Well drained soils, tolerates light frost, grows well in flood plains and swamps, stands dry period upto 7 months, precipitation: 200-1000 mm/yr, 0-40 °C and elevation upto 700 m	Fuel, poles and fence posts, charcoal, shelterbelt, apiculture & tool handles.	0.89		5-10 m <sup>3</sup> /ha/yr
13 <i>Eucalyptus tereticornis</i>	Sufeda, Lachi, Mysore hybrid	Myrtaceae	Australia	Plains and hills.	Well drained soils including saline/sodic/waterlogged soils, frost hardy and drought tolerant, precipitation: 800-1500 mm/yr, 0-40 °C, elevation: 1500 m.	Fuel, carriages, shelterbelt, furniture, fiber board, erosion control	0.7	4900 kcal/kg	12-25 m <sup>3</sup> /ha/yr
14 <i>Leucaena leucocephala</i>	Subabul, Ipil Ipil	Leguminosae- Mimosoideae	Mexico	Plains and foothills.	Tolerant, adaptable growing even on steep and marginal lands, grows in saline and acidic soils, precipitation: 500-1000 mm/yr, 2-45 °C and elevation upto 500 m.	Poles and construction, fuel, fodder, apiculture, agricultural implements, furniture and soil stabilization.	0.56	4600 kcal/kg	30 m <sup>3</sup> /ha/yr

No. Scientific name	Local name	Family-Subfamily	Origin	Distribution	Habitat & Ecology	Uses	Specific Gravity	Calorific Value	Productivity
15 <i>Mangifera indica</i>	Aam, Mango	Anacardiaceae	Pakistan, India, Nepal, Bhutan	River valleys of Chenab and Ravi, and in irrigated lands in Sindh.	Moderately shade tolerant, variety of well drained soils but does best on deep loamy soil, adaptable, humid hot/subtropical to tropical/monsoon climate with precipitation: 750-1500 mm/yr, -3.5 to 40 °C, elevation upto 600m	Fruit, chipboard, lumber and construction, ornamental, food and medicinal.	0.55	4600 kcal/kg	average height for 7 years is 4 meter
16 <i>Melia azedarach</i>	Bakain, Persian Lilac	Meliaceae	Lower Himalayas (Pakistan and Nepal)	Plains of Punjab and KPK	Well drained soils of valleys and ravines, mature tree is frost and drought hardy, easily coppiced, tropical to subtropical temperate climate, precipitation 600-1000 mm/yr, -5 -40 °C, elevation range of 900-1700 m.	Veneer and plywood, construction, boxes and crates, furniture, timber, agricultural implements, sports equipment, fodder, ornamental and medicinal.	0.56	5100 kcal/kg	17.5 m <sup>3</sup> /ha/yr
17 <i>Moringa oleifera</i>	Sohanjna, Horseradish Tree	Moringeaceae	Sub-Himalayan tract in Pakistan	Plains eastward from Rawalpindi	Intolerant, variety of soils does best in well drained soil, easily coppiced, endures drought periods and can grow on eroded sites.	Ornamental, food, fodder, seed oil, gum and medicinal.			

No. Scientific name	Local name	Family-Subfamily	Origin	Distribution	Habitat & Ecology	Uses	Specific Gravity	Calorific Value	Productivity
18 <i>Morus alba</i>	Tut, Mulberry	Moraceae	Pakistan, China, Central Asia, Afghanistan		Moderately intolerant, variety of well drained rich soils, semi-arid cool/cold subalpine/ subtropical winter/monsoon climate, -10-40 °C, precipitation: 750-1250 mm/yr, elevation upto 3300 m, frost hardy and drought tolerant if irrigated	Silkworm food, fodder, fruit, shelterbelts, carriages, sports goods, veneer, plywood, furniture and medicinal.	0.69	5100 kcal/kg	8.5 m <sup>3</sup> /ha/yr
19 <i>Populus deltoides</i>	Sufed Poplar, Northern Cottonwood	Salicaceae	North America	Plains and hills.	Sandy loams/alluvial soils, semi humid to semi arid/cool subtropical temperate climate, precipitation: 750-1250 mm/yr, -20-35 °C	Fuel, crates, packing cases, matches, pulp, reforestation, plywood, erosion control, fodder and roadside tree.	0.46	5900 kcal/kg	20-40 m <sup>3</sup> /ha/yr
20 <i>Populus euphratica</i>	Bahan, Euphrates Poplar	Salicaceae	Middle East, Southern Russia, Subcontinent, East to China	Hot arid areas with river sources/sub surface water	Variety of sites including waterlogged/saline soils, arid/ semi arid/ sub tropical climate, -10-45 °C, precipitation: 750-1250 mm/yr, elevation: upto 4000 m, frost hardy, drought resistant and withstands inundation, also considered a riverain species	Timber, packing cases, crates, matches, erosion control, reforestation, plywood, pulp, fodder, roadside tree and medicines	0.46	5000 kcal/kg	8-15 m <sup>3</sup> /ha/yr

No. Scientific name	Local name	Family-Subfamily	Origin	Distribution	Habitat & Ecology	Uses	Specific Gravity	Calorific Value	Productivity
21 <i>Populus euramericana</i>	Doghla Poplar, Hybrid Poplar	Salicaceae	Hybrid cultivar		Intolerant, deep soils with abundant water, withstands freezing temperatures but does not tolerate temperature exceeding 40-45 °C	Timber, packing cases, crates, matches, plywood, pulp, fodder, fuel, housing, furniture, chip board and shuttering poles.	0.28-0.52	5000 kcal/kg	40 m <sup>3</sup> /ha/yr
22 <i>Populus nigra</i>	Siah Poplar, Lombardy Poplar	Salicaceae	Western and Central Europe, Subcontinent, Middle East	Northern areas, Azad Kashmir, Plains throughout the country	Well drained soils along water courses, arid/cool-cold/subtropical climate, 20-45 °C, precipitation: 650-900 mm/yr, elevation range 900-3750 m, frost hardy	Fuel, packing cases, crates, matches, erosion control, reforestation, plywood, pulp, fodder, roadside plantation, general construction.	0.46	5000 kcal/kg	10-15 m <sup>3</sup> /ha/yr
23 <i>Prosopis cineraria</i>	Jand, Kandi	Leguminosae-Mimosoideae	Pakistan, India, Afghanistan, Middle East	Dry plains/hills of Sindh, Punjab, KPK and Balochistan	Variety of dry sites mostly clays/sands, does well in high alkaline soil, hot arid/semi arid to subtropical climate, -6-45 °C, precipitation: 75-650 mm/yr, elevation upto 450 m	Fodder, fuel, poles and construction, agriculture implements, apiculture, furniture, soil stabilization.	0.61	5000 kcal/kg	3-5 m <sup>3</sup> /ha/yr

No. Scientific name	Local name	Family-Subfamily	Origin	Distribution	Habitat & Ecology	Uses	Specific Gravity	Calorific Value	Productivity
24 <i>Prosopis juliflora</i>	Mesquite	Leguminosae-Mimosoideae	West Indies, Central and South America, Southwestern United States.	Dry plains/hills of Sindh, Punjab, KPK and Balochistan	Variety of dry sites mostly clays to sands, does well on high alkaline sites, drought and frost hardy, -2-45 °C, precipitation: 150-750 mm/yr, elevation upto 1200 m. Large tap root and extensive root system.	Fodder, fuel, poles and construction, agriculture implements, apiculture, furniture, soil stabilization.	0.7	4500 kcal/kg	3-5 m <sup>3</sup> /ha/yr
25 <i>Syzygium cumini</i>	Jamun, Jaman, Black Plum	Myrtaceae	Subcontinent	Plains and lower hills of Punjab, KPK and Azad Kashmir.	Well drained soils from sands to loams, semi humid warm hot/subtropical winter/monsoon climate, -5-40 °C, precipitation: 1250 mm/yr, and elevations upto 1500 m, frost hardy and if irrigated then tolerates drought	Construction, fruit, fuel, paper pulp, apiculture, medicinal, tannin, shelterbelts, roadside planting and shade.	0.7	4800 kcal/kg	12 m <sup>3</sup> /ha/yr
26 <i>Tamarindus indica</i>	Imli, Tamarind	Leguminosae-Caesalpinioideae	Tropical Africa	Sindh, Punjab	Variety of soils but does best in deep alluvium soils and dry sandy sites as well, hot humid/dry tropical to subtropical climate, 0-40 °C, precipitation: 250-1250 mm/yr, elevation upto 600 m	Implement handles, fuel, charcoal, furniture, wheels, axles, food, fodder, apiculture, ornamental and medicinal.	0.91-1.28	4969 kcal/kg	height increased @ 0.8m/yr

No. Scientific name	Local name	Family-Subfamily	Origin	Distribution	Habitat & Ecology	Uses	Specific Gravity	Calorific Value	Productivity
27 Tamarix aphylla	Frash, Ghaz, Khaggal, Tamarisk	Tamaricaceae	Middle East, Pakistan, Central Asia, North Africa	Plains of Punjab, KPK, Sindh, Balochistan and Thal area.	Well drained sandy soils including saline/sodic sites, arid o hot subtropical winter monsoon climate, - 1-50 °C, precipitation: 100-500 mm/yr, does not grow on elevation of more than 600 m, frost hardy	Carpentry, agriculture implements, fuelwood, shelterbelts, charcoal, tannin, erosion control, sand dune stabilization.	0.68	4835 kcal/kg	5-10 m <sup>3</sup> /ha/yr
28 Terminalia arjuna	Arjun	Combretaceae	Subcontinent	Plains, gardens, roadside plantation	Shade tolerant, variety of moist well drained sites, also grows on saline/sodic/waterlogged soils, humid hot tropical/subtropical monsoon climate, 0-45 °C, precipitation: 750-3800 mm/yr, elevation: 600m.	Fuel, implements, erosion control, wheels, fodder, ornamental, timber and medicines.	0.9	5000 kcal/kg	10-12 m <sup>3</sup> /ha/yr
29 Zizyphus mauritiana	Ber, Chinese Date	Rhamnaceae	South Asia	Throughout the country	No particular soil requirement, warm temperate to subtropical to tropical climate, -5-50 °C, precipitation 600-1500 mm/yr, elevation upto 600m.	Fuel, charcoal, agricultural implements, fruit.	0.93	5900 kcal/kg	average height for 7 years is 5 meter

## 9.10. Conclusion

Agroforestry is the purposeful growing or deliberate retention of trees with crops and/or animals in interacting combinations for multiple products or benefits from the same management unit. Furthermore, it is classified in three main classes or agroforestry systems i.e. agrisilviculture, silvipastoral and agrisilvopastoral systems. Poor soils adversely affect the biomass production and yield of agricultural crops. Large area of the world has been classified as degraded soils and bringing such area under cultivation is a big challenge for farmers. However, promotion of agroforestry practices on these areas can provide manifold benefits to farmers like soil reclamation, improving soil fertility through nitrogen fixation and increasing tree cover. Moreover, planting multipurpose trees can also fulfill food and fuelwood requirements of local inhabitants/ farmers.

## References

- Allen, S.C., S. Jose, P.K.R. Nair, B.J. Brecke, P. Nkedi-Kizza and C.L. Ramsey (2004). Safety-net role of tree roots: Experimental evidence from an alley cropping system. *For. Ecol. Manage.* 192: 395-407.
- Ansari, M.A.A. and M. Iftikhar (1985). Food and forest. *Pak. Agric.* 7: 25-29.
- Bhardwaj, S.P. and O.P.S. Khola (1999). Biological and mechanical measures for erosion control and crop production on 4 percent slope. Annual Report, CSWSRTI, Dehradun, India, pp. 22.
- Bhardwaj, S.P. (1990). Vegetative barrier is an effective economic and eco-friendly measure for erosion control on agricultural lands. In: Abstracts of 8<sup>th</sup> ISCO conference, New Delhi, India. pp. 203-206.
- Blanford, H.R. (1958). Highlights of one hundred years of forestry in Burma. *Empire Fores. Rev.* 37: 33-42.
- Cavalcante, A.M.B. and S.C.J.G. de A. Perez (1995). Efeitos dos estresses hídrico e salino sobre a germinação de sementes de *Leucaena leucocephala* (Lam.) de Witt. [Effects of water and salt stress on the germination of seeds of *Leucaenaleucocephala* (Lam.) de Witt]. *Pesquisa Agropecuária Brasileira.* 30: 281-289.
- FAO (2000). Global Forest Resources Assessment 2000 and UNEP Global Biodiversity Outlook 2001 [Online] <http://www.safnet.org/aboutforestry/world.cfm>, Accessed on April 12, 2014.
- FAO (2008). The state of food and agriculture. Biofuels: prospects, risks and opportunities. Food and Agriculture Organization of the United Nation, Rome, Italy.
- FAO (2010). Global forest resources assessment 2010. FAO Forestry Paper No. 163. Food and Agriculture Organization of the United Nation, Rome, Italy.
- Fischer, M., K. Cruickshanks, B. Klein, R. Klein, C.R. Schubert and T.L. Wiley (2009). Multiple sensory impairment and quality of life. *Ophthalmic Epidemiol.* 166: 346-353.
- Flowers, T. J. (2004). Improving crop salt tolerance. *J. Exp. Bot.* 55: 307-319.

- Gogoi, B.N. (1976). A review of research in shade tea in India. Two and a bud. 23: 67-73.
- Government of Pakistan (1992). Forestry Sector Master Plan, Ministry of Planning and Development in Collaboration with Asian development Bank and United Nations Development Program: I-III, [Online] <http://www.pakistan.gov.pk>, Accessed on July 26, 2014.
- Government of Pakistan (2005). Economic Survey of Pakistan, Economic Advisor's Wing, Finance Division, Islamabad, Pakistan, pp.11-26.
- Government of Pakistan (2011). Pakistan: new growth framework, Government of Pakistan, Planning Commission, Islamabad., 2011, pp-33.
- Government of Pakistan (2014). Economic Survey of Pakistan, Economic Affairs Division, Ministry of Finance, Islamabad, Pakistan.
- Grewal, S.S. (1993). Agroforestry systems for soil and water conservation in Shivaliks. In: Agroforestry in 2000 AD for the semi-arid and arid tropics, pp: 82-85, National Research Center for Agroforestry, Jhansi, India.
- Hartemink, A.E., R.J. Buresh, B. Jama and B.H. Janssen (1996). Soil nitrate and water dynamics in sesbania fallows, weed fallows and maize. Soil Sci. Soc. Amer. J. 6: 568-574.
- ICFRE (1993). Annual Research Report, 1991-92. Indian Council of Forestry Research and Education, Dehradun, India. pp. 81.
- Jain, S.K. and P. Singh (2000). Economic analysis of industrial agroforestry: poplar (*Populus deltoids*) in Uttar Pradesh (India). Agrofor. Syst. 49: 255-273.
- Jamnadass, R.H, I.K. Dawson, S. Franzel, R.R.B. Leakey, D. Mithöfer, F.K. Akinnifesi and Z. Tchoundjeu (2011). Improving livelihoods and nutrition in sub-Saharan Africa through the promotion of indigenous and exotic fruit production in smallholders' agroforestry systems: a review. Int.For.Rev.13: 338-354.
- Kort, J. (1988). Benefits of windbreaks to field and forage crops. Agric. Ecosys. Environ. 22/23: 165-190.
- Krohn, T.J. (1981). Fertilization of *Eucalyptus tereticornis* in Southern Guam. Tree Planters' Notes 32(1): 26-28.
- Leakey, R.R.B. (1999). Potential for novel food products from agroforestry trees. Food Chem.64: 1-14.
- Malézieux, E. (2013). Underutilized fruit trees in Africa. Special Issue. *Revue Fruits* (in press).
- Maslekar, A.R. (1990). Foresters Companions. Jugal Kishore and Co. (Publication Division), Dehradun, India. pp. 603.
- McAdam, J.H., T.H. Thomas and R.W. Willis (1999). The economics of agroforestry systems in the United Kingdom and their future prospects. Scottish Fores. 53: 37-41.
- Mercer, D.E. and R.P. Miller (1998). Socioeconomic research in agroforestry: progress, prospects, priorities. Agrofor.Syst. 38: 177-193.
- Miiller, E.U. and S.J. Scherr (1990). Planning technical interventions in agroforestry projects. Agrofor.Syst.11: 23-44.
- Mohsin, F., R.P. Singh and K. Singh (1996). Nutrient cycling of poplar plantation in relation to stand age in agroforestry system. Ind. J. Fores. 19: 302-310.

- Munns, R. (2002). Comparative physiology of salt and water stress. *Plant Cell Environ.* 25: 239-250.
- Munns, R. and M. Tester (2008). Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* 59: 651-681.
- Nair, P.K.R. (1985). Classification of agroforestry systems. *Agrofor. Syst.*3: 97-128.
- Nair, P.K.R. (1987). Agroforestry systems inventory. *Agrofor. Syst.*5: 301-317.
- Nair, P.K.R. (1992). State-of-the-art of agroforestry research and education. Keynote paper to the International Conference on Agroforestry for Sustainable Development. Univ. of Chapingo, Texcoco, Mexico.
- Nair, P.K.R. (1993). The History of Agroforestry. In: An Introduction to Agroforestry. Klgwer Academic Publishers, International Center for Research in Agroforestry, Nairobi, Kenya. pp. 3-11.
- Nair, P.K.R. (1996). Agroforestry directions and literature trend. In: McDonald, P. and J. Lassoie (ed). *The Literature of Forestry and Agroforestry*. Cornell University Press, USA. pp: 74-95.
- Narain, P., R. Singh and P. Joshi (1988). Studies on nutrient and water budget under different land use systems. Annual Report, pp: 55-57, Central Soil and Water Conservation Research & Training Institute, Dehradun, India.
- Quraishi, M.A.A. (1998). *Basics of Forestry and Allied Sciences*. Vol. 1, A-One Publishers, Lahore, Pakistan. pp. 86-92.
- Raghavan, M.S. (1960). Genesis and history of the Kumri system of cultivation. In: *Proceedings of the Ninth Silviculture Conference*, Dehra Dun, 1956. Forest Research Institute, Dehra Dun, India.
- Rai, S.N. (1996). Prominent findings of forestry research in India. FORSPA Publication., Bangkok, Thailand. pp. 79.
- Ramadhani, T., R. Otsyina and S. Franzel (2002). Improving household incomes and reducing deforestation; the example of rotational woodlots in Tabora District, Tanzania. *Agric. Ecosys. Environ.* 89: 227-237.
- Samra, J.S. (1994). Strategies for torrent control in Shiwaliks. *Indian J. Soil Conserv.* 22: 134-144.
- Scherr, S.J. (2004). Building opportunities for small-farm agroforestry to supply domestic wood markets in developing countries. *Agrofor. Syst.* 61: 357-370.
- Sen, W. (1991). *Agroforestry in China*. Ministry of Foreign Affairs, Beijing, China.
- Sileshi, G.W., F.K. Akinnifesi, O.C. Ajayi and B. Muys (2011). Integration of legume trees in maize based cropping systems improves rain-use efficiency and yield stability under rain-fed agriculture. *Agric. Water Manage.*98: 1364-1372.
- Sileshi, G.W., L.K. Debusho and F.K. Akinnifesi (2012). Can integration of legume trees increase yield stability in rainfed maize cropping systems in Southern Africa? *Agron. J.*104: 1392-1398.
- Singh, G., G. Sastry and S.P. Bhardwaj (1990). Watershed responses to conservation measures under different climatic regions of India. *Indian J. Soil Conserv.* 18: 16-22.
- Singh, G., R. Babu and S. Chandra (1981). Soil loss prediction research in India. ICAR Bulletin T-12/D-9. Central Soil and Water Conservation Research and Training Institute, Dehra Dun, India. pp. 70.

- Soemarwoto, O. and G.R. Conway (1991). The Javanese Homegardens. *J. Farm. Sys. Res. Ext.* 2: 95-118.
- Sreemannarayana, B., L.G.G. Rao and B. Joseph (1994). Evaluation of multipurpose tree species and their influence on soil fertility improvement. *International conference: special issue. Range Manage. Agrofor.* 15: 199-202.
- Szott, L.T., C.A. Palm and P.A. Sanchez (1991). Agroforestry in acid soils in the humid tropics. *Adv. Agron.* 45: 275-301.
- Tejwani, J.G (1994). *Agroforestry in India*. Oxford & IBH Publishing Co Pvt. Ltd., New Delhi, India. pp. 233.
- Thorlakson, T. and H. Neufeldt (2012). Reducing subsistence farmers' vulnerability to climate change: evaluating the potential contributions of agroforestry in western Kenya. *Agric.Food Secur.* 1:15.
- UN. (1995). United Nations (U.N.) Population Division, *World urbanization Prospects: The 1994 Revision*. New York, USA. pp. 87.
- Wani, B.A. (2003). National Report to the Third Session of the United Nations Forum on Forests (UNFF): 1-7, [Online] `Accessed on January 06, 2014.
- Williams, P.A., A.M. Gordon, H.E. Garrett and L. Buck (1997). Agroforestry in North America and its role in farming systems. In: Gordon, A.M. and S.M. Newman (ed). *Temperate Agroforestry Systems*. CABI, Wallingford, UK. pp. 9–84.
- World Bank. (2011). *Wood-based biomass energy development for sub-Saharan Africa: issues and approaches*. Washington, DC.
- Wu, Y and Z. Zhu (1997). Temperate agroforestry in China. In: Gordon, A.M. and S.M. Newman (ed). *Temperate Agroforestry Systems*. CABI, Wallingford, UK. pp. 149–179.
- Young, A. (1997). *Agroforestry for soil management*. CAB Int.Wallingford, UK. pp. 320.
- Young, A. (1986). The potential of agroforestry for soil conservation and sustainable land use. In: Kozub, J. (ed). *Land and Water Resources Management*. Washington, DC: Economic Development Institute of the World Bank. pp. 301-317.
- Zaigham, N.A. and A.N. Zeeshan (2004). Prospects of Renewable Energy Sources in Pakistan. In: Khan, H.A., M.M. Qureshi, T. Hussain and I. Hayee (ed). *Proceedings of COMSATS Conference 2004 on Renewable Energy Technologies & Sustainable Development*. Commission on Science and Technology for Sustainable Development in South G-5/2, Islamabad (Pakistan); 154 p
- Zaman, S.B. and S. Ahmad (2012). Wood supply and demand analysis in Pakistan – key issues, *Managing Natural Resources for sustaining Future Agriculture*, Research Briefings: 4(22):1-12. Pakistn Research Council Islamabad, Pakistan.
- Zhu, J.K., P.M. Hasegawa and R.A. Bressan (1997). Molecular aspects of osmotic stress. *Crit. Rev. Plant Sci.* 16: 253-277.